

RESEARCH STATEMENT

ZACH TEITLER

My area of research is commutative algebra and algebraic geometry (MSC 13 and 14).

Together with a number of collaborators, I have worked on broad range of problems, often with a combinatorial or computational flavor. These projects have involved a variety of topics including secant varieties and Waring rank, multiplier ideals, computational experimentation, arrangements of points and hyperplanes, and more. I look forward to continuing this work, as well as pursuing future collaborations in new directions.

1. WARING RANK

The *Waring rank* of a homogeneous form $F = F(x_1, \dots, x_n)$ of degree d is the least r such that $F = c_1 \ell_1^d + \dots + c_r \ell_r^d$ for some linear forms ℓ_i and scalars c_i . For example,

$$xy = \frac{1}{4}(x+y)^2 - \frac{1}{4}(x-y)^2,$$

so the rank $r(xy) \leq 2$; and $r(xy) \geq 2$ as xy is not equal to a square of a linear form.

Waring ranks and related questions have been studied since the mid-19th century (for example [67], see also [35, 43] and discussions of applications in [21, 47, 48]) but the last 5 years have seen explosive activity and rapid advances.

For example, Carlini, Catalisano, and Geramita have determined the Waring rank of a monomial [20]: $r(x_1^{a_1} \cdots x_n^{a_n}) = (a_1 + 1) \cdots (a_n + 1)$, when $0 < a_1 \leq a_2 \leq \dots \leq a_n$. It is remarkable that this was determined in 2011, after more than 150 years of study of Waring ranks. They also show that the Waring rank of a sum of monomials in separate variables is equal to the sum of the Waring ranks of the separate monomials. It is conjectured that, in general, $r(F(x) + G(y)) = r(F) + r(G)$; this remains open after decades of work, but building on the 2011 result for sums of monomials, progress has been made by Y. Woo [81] and Carlini–Catalisano–Chiantini [19].

The primary tools for studying Waring rank arise from commutative algebra. For $F \in S = \mathbb{C}[x_1, \dots, x_n]$, the **dual ring** is $T = \mathbb{C}[\partial_1, \dots, \partial_n]$, acting on S by having each ∂_i act as $\partial/\partial x_i$; this action is the **apolarity pairing**, giving a perfect pairing $T_d \times S_d \rightarrow \mathbb{C}$ for each d (and $T_a \times S_d \rightarrow S_{d-a}$ for all d, a). Equivalently, the commutative algebra notion of **catalecticant map** of a given form $F \in S_d$ is the linear map $C_F^a : T_a \rightarrow S_{d-a}$, for $0 \leq a \leq d$, sending differential operators to their evaluations on F .

My contributions include the following.

1.1. An upper bound for rank and generalized rank. Let $X \subset \mathbb{P}^N$ be a nondegenerate projective variety. The X -rank of a point q is the least number of points on X linearly spanning a space containing q . This includes Waring rank as the case X is a Veronese variety. It also includes ordinary tensor rank (see for example [47]) as the case X is a Segre variety. Many other natural notions of rank can be expressed as X -rank for an appropriately chosen X .

Grigoriy Blekherman and I have given the following upper bound for X -rank, for any irreducible nondegenerate variety X over a closed field in any characteristic:

Theorem 1 ([7]). *Let r_g be the generic rank with respect to X , i.e., the rank of a general point. Then for all $q \in \mathbb{P}^N$, the X -rank of q is at most $2r_g$.*

This is a significant improvement for a number of previously studied cases, most notably Waring rank. A further improvement is shown in the case that X has some higher secant variety of codimension 1. We also give the following result for real rank.

Theorem 2 ([7]). *Let $X \subset \mathbb{R}\mathbb{P}^N$ be a real nondegenerate projective variety such that the real points of X are dense in X . Let r_0 be the smallest typical real rank with respect to X and let r_g be the generic rank with respect to the complexification $X_{\mathbb{C}} = X \otimes \mathbb{C}$. Then $r_0 = r_g$ and for all $q \in \mathbb{R}\mathbb{P}^N$, the X -rank of q is at most $2r_0$.*

Future work. I will study homogeneous forms and tensors which are believed to have high rank, in order to narrow down the range of possible values for the maximum rank.

Previous upper bounds for Waring rank are asymptotically not as good as the above, but they still raise intriguing questions. For example, the papers [5, 44, 2] actually give an upper bound for “open Waring rank,” which is greater than ordinary Waring rank; I will investigate the value of open Waring rank for a general form.

1.2. Power sum decompositions of monomials. This is joint work with Weronika Buczyńska and Jarosław Buczyński.

Theorem 3 ([14]). *Let $F = x_1^{a_1} \cdots x_n^{a_n}$ be a monomial, $0 < a_1 \leq \cdots \leq a_n$, and let $r = r(F) = (a_2 + 1) \cdots (a_n + 1)$. Let ℓ_1, \dots, ℓ_r be linear forms and let $I \subset \mathbb{C}[y_1, \dots, y_n]$ be the homogeneous defining ideal of the set of projective points $\{[\ell_1], \dots, [\ell_r]\}$. Then $F = c_1 \ell_1^d + \cdots + c_r \ell_r^d$ for some scalars c_i if and only if I is a complete intersection of type $(a_1 + 1, \dots, a_n + 1)$, generated by*

$$y_1^{a_1+1} - \phi_1 y_0^{a_0+1}, \dots, y_n^{a_n+1} - \phi_n y_0^{a_0+1}$$

for some forms ϕ_i of degree $a_i - a_0$.

This theorem, together with some additional reductions, allows the computation of the dimension of the variety

$$(1) \quad \left\{ \{[\ell_1], \dots, [\ell_r]\} : F = c_1 \ell_1^d + \cdots + c_r \ell_r^d, \text{ some } c_i \right\}.$$

In particular let $(\mathbb{C}^*)^n$ act by scaling the variables and let $T \subset (\mathbb{C}^*)^n$ be the subtorus that fixes F .

Corollary 4 ([14]). *The induced action of T on the variety (1) is transitive if and only if $a_1 = \cdots = a_n$.*

Thus the minimum length power sum decomposition of a monomial is unique up to scaling variables if and only if the monomial is of the form $(x_1 \cdots x_n)^k$.

Future work. Carlini–Catalisano–Geramita, in addition to determining Waring ranks of monomials, showed that the Waring rank of a sum of pairwise coprime monomials (equivalently: monomials in independent variables) is equal to the sum of the Waring ranks of the separate monomials. (This is conjectured to hold for all sums of forms in independent variables, see [81, 19].) An obvious conjecture is that every Waring decomposition (= minimum length power sum decomposition) of a sum of pairwise coprime monomials must be given

by a concatenation of separate Waring decompositions of the separate monomials. This is open, even for 2-term sums.

I will study the generalization of the above theorem to other forms with complete intersection apolar annihilating ideals, such as defining equations of reflection arrangements of hyperplanes.

1.3. Sub-generality of ranks of monomials. Not many homogeneous forms are known to have higher than generic rank; it has not even been shown that forms with higher than generic rank exist in all degrees and numbers of variables (although it would be very surprising to learn otherwise).

Following the determination by Carlini–Catalisano–Geramita of the ranks of monomials and sums of pairwise coprime monomials, it is natural to ask if any of them have higher than generic rank. And indeed, Carlini–Catalisano–Geramita observed that infinitely many monomials in three variables have higher than generic rank, but at most finitely many in n variables do so, for each $n \geq 4$. But they did not determine how many such monomials there were, or give any examples.

This was completed in joint work with three undergraduate students Erik Holmes, Paul Plummer, and Jeremy Siegert.¹

Theorem 5 ([42]). *Every monomial in 4 or more variables has Waring rank strictly less than the generic Waring rank. Furthermore, in 4 or more variables, every homogeneous sum of pairwise coprime monomials has Waring rank strictly less than the generic Waring rank, with exactly three exceptions: $x_1x_2^2+x_3x_4^2$ has Waring rank 6, strictly greater than the generic Waring rank of forms of degree 3 in 4 variables, which is 5; $x_1x_2^2+x_3^3+x_4^3$ has Waring rank 5 and $x_1x_2x_3+x_4^2$ has Waring rank 5, equal to the generic Waring rank.*

I proved the statement on monomials and conjectured the statement on sums of monomials, which was then proved by the three undergraduates.

Future work. It is not known what is the maximum Waring rank attained by sums of pairwise coprime monomials (of a given degree, in a given number of variables); the result above is shown by giving an upper bound which is less than the generic Waring rank, but the upper bound is not actually attained. A number of examples found by the three undergraduate students show that the rank of such a sum is not maximized by making “greedy” choices, i.e., choosing each term to be the monomial of largest rank possible with the number of remaining variables.

This is a suitable question for a group of undergraduate students. Small cases should be amenable to computer exploration. It will be a good opportunity for new researchers to use a combination of computer programming and exhaustive search, pattern recognition, and looking for a proof idea.

1.4. A lower bound for ranks of invariant forms. This is joint work with Harm Derksen. In the following statement, for any polynomial P , let $\text{Diff}(P)$ be the vector spanned by the derivatives of P of all orders, including P itself (zeroth order derivative):

¹Erik Holmes graduated from Boise State in 2014 and began the Ph.D. program at University of Hawaii. Paul Plummer graduated from Boise State in 2014 as well and is currently applying for Ph.D. programs. Jeremy Siegert will graduate from Boise State in December 2014 and is also applying for Ph.D. programs.

Theorem 6 ([25]). *Let G be a connected group with an irreducible representation V and let F be an invariant form on V . Then the Waring rank of F is bounded below by*

$$\dim \text{Diff}(F) - \dim \text{Diff}(\partial F / \partial x)$$

for any nonzero $x \in V$.

More generally, the main result of our paper gives a lower bound for simultaneous Waring rank of an invariant linear series of forms, and allows reducible representations (then, roughly speaking, $x \in V$ must simply be chosen not lying in any proper subrepresentations). The same result is shown to hold for arbitrary forms (or linear series), not necessarily invariant under any group action, when $x \in V$ is general. These bounds are actually lower bounds for cactus rank, which is the following modification of Waring rank. The cactus rank $cr(F)$ is the least r for which there exists a zero-dimensional scheme Z of length r such that F lies in the linear span of the degree d Veronese re-embedding of Z . Waring rank is the case when Z is also required to be reduced, $Z = \{[\ell_1], \dots, [\ell_r]\}$.

For example, the generic $n \times n$ determinant $\det_n = \det((x_{i,j})_{1 \leq i,j \leq n})$ is an invariant form under left and right multiplication by SL_n and we get $r(\det_n) \geq \binom{2n}{n} - \binom{2n-2}{n-1}$. This is the best currently known lower bound for rank of the determinant.

Future work. I will study apolarity and Waring ranks of forms invariant under finite groups, such as symmetric polynomials, and also forms invariant under disconnected groups, such as the generic permanent. The permanent in particular is related to a version of P versus NP in geometric complexity theory.

I will also study upper bounds for the rank of determinant, building on previous work of Derksen.

1.5. Reflection multiarrangements. Alex Woo and I determined the Waring rank of many defining equations of reflection multiarrangements, i.e., fundamental skew invariants of finite complex reflection groups.

Theorem 7 ([75]). *Let G be a finite complex reflection group on \mathbb{C}^n with degrees $1 \leq d_1 \leq \dots \leq d_n$. We do not assume G acts essentially. Let f_G be the defining equation of the reflection arrangement of G (equivalently: the fundamental skew invariant of G , or the Jacobian of the fundamental invariants of G). Then the cactus rank $cr(f_G)$ is*

$$cr(f_G) = \frac{|G|}{d_n} = d_1 \cdots d_{n-1}$$

and the Waring rank $r(f_G)$ is bounded by

$$\frac{|G|}{d_n} \leq r(f_G) \leq \frac{|G|}{D},$$

where D is the greatest regular number of G . The upper bound for Waring rank is given by an explicit power sum decomposition.

Note that $d_n = D$ is a regular number for many reflection groups of interest, including all irreducible real reflection groups and many irreducible complex reflection groups. For example, the fundamental skew invariant of the symmetric group S_n is the classical Vandermonde determinant

$$f_{S_n} = \prod_{i < j} x_j - x_i,$$

and we obtain $r(f_{S_n}) = (n - 1)!$.

Future work. I will study other complex reflection groups. Perhaps the simplest example of a complex reflection group whose greatest degree is not a regular number is the group $G = \mathbb{Z}/a_1\mathbb{Z} \times \cdots \times \mathbb{Z}/a_n\mathbb{Z}$ with not all a_i equal. In this case the fundamental skew invariant is the monomial $x_1^{a_1} \cdots x_n^{a_n}$, whose rank was determined by Carlini–Catalisano–Geramita using commutative algebra. The same algebraic technique should be tried for other examples.

For instance, one may consider the reducible reflection group given by a product of symmetric groups $G = S_{a_1} \times \cdots \times S_{a_n}$ with not all a_i equal. In this case the fundamental skew invariant is a product of classical Vandermonde determinants in separate variables.

I will work with Stefan Tohaneanu and Alex Woo to describe the apolar annihilating ideals of general hyperplane arrangements, in particular the Hilbert functions, graded Betti numbers, and minimal generators of those ideals, and bound or determine the Waring ranks of such arrangements. It might be possible to also describe the minimal free resolution of the annihilating ideals but this is more speculative. At this point we are able to produce candidate generators of the ideal via Proposition 3.3 of [73] which relates apolarity of F to singularities of the hypersurface $V(F)$: our candidate generators correspond naturally to the 0-skeleton of the (projectivized) hyperplane arrangement.

1.6. Apolarity and direct sum decompositions. In joint work with Weronika Buczyńska, Jarosław Buczyński, and Johannes Kleppe, the theory of apolarity is used to study decompositions of a form F as a sum of forms depending on linearly independent sets of variables, possibly after a linear change of coordinates. For example, it is clear that $xy \neq G(x) + H(y)$, but $xy = G(\ell_1) + H(\ell_2)$, namely,

$$xy = \frac{1}{4}\ell_1^2 - \frac{1}{4}\ell_2^2$$

for $\ell_1 = x + y$, $\ell_2 = x - y$. In general, for $F = F(x_1, \dots, x_n)$, we ask whether there is an expression of the form

$$F = G(\ell_1, \dots, \ell_k) + H(\ell_{k+1}, \dots, \ell_n)$$

with the ℓ_i linearly independent linear forms. Such an expression is called a **direct sum decomposition** of F . For another example, consider the generic determinant $\det_n = \det((x_{i,j})_{1 \leq i,j \leq n})$. Then $\det_2 = x_{1,1}x_{2,2} - x_{1,2}x_{2,1}$ is visibly decomposed as a sum of forms in separate variables. It is easy to see that for $n > 2$, $\det_n \neq G(x_{1,1}, \dots) + H(\dots, x_{n,n})$, but it is less obvious whether after a linear change of coordinates (in n^2 variables!) \det_n is so decomposable. We find a negative answer:

Theorem 8 ([13]). *Let $F \in \mathbb{C}[x_1, \dots, x_n]$ be a homogeneous form of degree d . Let $F^\perp \subset T = \mathbb{C}[\partial_1, \dots, \partial_n]$ be the homogeneous ideal of differential operators D such that $DF = 0$.*

- (1) *If F is decomposable as a direct sum then F^\perp has a minimal generator of degree d .*
- (2) *If F^\perp has a minimal generator of degree d then F is a limit of forms which are decomposable as direct sums.*
- (3) *Suppose that F cannot be written as a form in fewer variables, even after a linear change of coordinates. Then F is a limit of forms which are decomposable as direct sums if and only if F^\perp has a minimal generator of degree d .*

In particular, due to Shafiei's proof of my conjecture that $(\det_n)^\perp$ is generated in degree 2 [61], while \det_n has degree n , it follows that \det_n is indecomposable as a direct sum for $n > 2$. Shafiei's further results regarding permanents, pfaffians, symmetric determinants,

symmetric permanents, etc., have similar implications for indecomposability of those forms [61, 60].

Many more results are given in [13], such as a lower bound on the greatest degree of a minimal generator of F^\perp .

Theorem 9 ([13]). *Let F be a form of degree d in n variables and suppose that F^\perp is generated in degrees less than or equal to δ . Then $d \leq (\delta - 1)n$.*

Future work. I will study generalizations to local rings (such as formal power series) in place of polynomial rings. I will also study forms that make the lower bound sharp, $d = (\delta - 1)n$.

1.7. A lower bound for Waring rank. This is joint work with J.M. Landsberg. We give a geometric lower bound for Waring rank, specifically a lower bound for $r(F)$ in terms of the singularities of the hypersurface defined by F .

Theorem 10 ([49]). *Let F be a form of degree d . Fix an integer $0 < a < d$. Let C_F^a be the a th catalecticant map of F . Suppose F cannot be written as a form in fewer variables, even after a linear change of coordinates (this is equivalent to the easily checked condition that C_F^1 be injective). Let $\Sigma_a(F)$ be the projective variety defined by the vanishing of the forms in the image of C_F^a , i.e., the set of a -th derivatives of F , so $\Sigma_a(F)$ is the variety of points at which F vanishes with multiplicity at least $a + 1$. Then $r(F) > \text{rank } C_F^a + \dim \Sigma_a(F)$.*

This is notable for being the only currently known lower bound for Waring rank which is not actually a lower bound for cactus rank.

Future work. Examples in [49] suggest approaches to search for improvements to the above bound, by incorporating more information about the singularities, beyond just the dimension of the singular locus.

1.8. Geometric lower bounds for generalized rank. I have generalized the lower bound for Waring rank (and cactus rank) discovered by Ranestad and Schreyer [58], and the lower bound for Waring rank discovered by Landsberg and myself [49], to lower bounds for ranks of linear series (simultaneous Waring rank), ranks of multihomogeneous forms, and ranks with respect to nondegenerate projective varieties X [73]. As a simple example, it is shown that there are linear forms ℓ_i and m_i , and scalars c_i , such that

$$x_1 \cdots x_a y_1 \cdots y_b = \sum_{i=1}^r c_i \ell_i(x)^a m_i(y)^b$$

if and only if $r \geq 2^{a+b-2}$.

Future work. The generalizations to X -rank are incomplete; a number of questions are posed in [73]. For example, for a form F in n variables and any $1 \leq k \leq n$ let $r_k(F)$ be the least number of terms in an expression for F as a sum of forms each depending on k or fewer variables (possibly after a linear change of coordinates; i.e., the extension of a form on a k -dimensional subspace), see [17, 18]. Waring rank is the case $k = 1$. If $0 < d_1 \leq \cdots \leq d_n$, then


$$(d_1 + 1) \cdots (d_{n-k} + 1) \leq r_k(x_1^{d_1} \cdots x_n^{d_n}) \leq (d_2 + 1) \cdots (d_{n-k+1} + 1)$$

and the right hand inequality is conjectured to be equality, see [73, Conjecture 5.18].

I will study and hopefully determine r_k of a monomial. This is very close to the determination of classical Waring rank of a monomial. The commutative algebra techniques used by Carlini–Catalisano–Geramita for that problem should be applicable here.

Many, many more questions are suggested in [73], including a number of questions that would be good projects for students.

1.9. **Interlude.** The power sum decomposition $xy = \frac{1}{4}(x+y)^2 - \frac{1}{4}(x-y)^2$ can be rewritten as $(x+y)^2 = 4xy + (x-y)^2$, and indeed, for $x > y > 0$, four $x \times y$ rectangles and an $(x-y) \times (x-y)$ square can be fitted together to form an $(x+y) \times (x+y)$ square:



Now $r(xyz) = 4$ and a power sum decomposition of length 4 is given by

$$xyz = \frac{1}{24} \left((x+y+z)^3 - (x+y-z)^3 - (x-y+z)^3 - (-x+y+z)^3 \right),$$

which can be rearranged to

$$(x+y+z)^3 = 24xyz + (x+y-z)^3 + (x-y+z)^3 + (-x+y+z)^3.$$

So it is natural to ask: if $x, y, z > 0$ are the sides of a triangle, then can $24 x \times y \times z$ “bricks”, plus three cubes as indicated, fit together to form a cube of side $x + y + z$?

Following Igor Pak’s suggestion, I obtained 24 bricks and 3 cubes of appropriate sizes created using a 3D printer, and engaged in experimental mathematics. I was able to show that the answer is *no* — for general x, y, z there is no such stacking. A simple counterexample is given by $(x, y, z) = (11, 13, 17)$. (For these values, I showed that each face of the big cube can only be covered by having exactly one small cube touching it, and the small cube must be centered on the face of the big cube; but there are 6 faces and only 3 small cubes, so it is impossible.)

2. MULTIPLIER IDEALS

Multiplier ideals have been applied to a number of problems in algebraic geometry in recent years, most spectacularly in recent major advances in the minimal model program [37, 6] that built on earlier work showing the deformation invariance of plurigena [63]. Other applications include several results on singularities and linear series [50], [31], a bound for symbolic powers [30], and applications to algebraic statistics [80], [82], [26, Chapter 5]. New applications of multiplier ideals continue to emerge in topics such as Chow stability [51] and singularities in generic liaison [57]. With broad and growing interest in multiplier ideals, it is increasingly valuable to compute examples.

For a thorough introduction to multiplier ideals see [50]. There are a number of equivalent characterizations of multiplier ideals, in terms of jet spaces [29], D -modules [16], test ideals for tight closure [64], and local integrability [45, 46, 55]. Here is a definition of multiplier ideals in terms of resolution of singularities. Suppose X is a smooth variety over a field

\mathbb{k} (we may assume X is affine, or even just \mathbb{k}^n , since we are primarily interested in local issues), $I \subset \mathcal{O}_X$ is a nonzero ideal sheaf, and $\mu : Y \rightarrow X$ is a log resolution of I , so that the total transform $I\mathcal{O}_Y$ defines a divisor F with simple normal crossings support, $F = \sum a_i E_i$, where the E_i are distinct reduced components of F . Then for each real number $c \geq 0$, the c 'th multiplier ideal is defined by $\mathcal{J}(I^c) = \mu_* \mathcal{O}_Y(K_{Y/X} - \lfloor c \cdot F \rfloor)$ where $K_{Y/X}$ is the relative canonical divisor of Y over X , defined locally by the vanishing of the determinant of the Jacobian $d\mu$, and $\lfloor c \cdot F \rfloor$ denotes the component-wise round-down of the \mathbb{R} -divisor $c \cdot F$, given by $\lfloor c \cdot F \rfloor = \sum \lfloor ca_i \rfloor E_i$.

My contributions include the following.

2.1. Multiplier ideals of hyperplane arrangements. Mustaă computed multiplier ideals of hyperplane arrangements using jet schemes [54]. I then gave a proof via resolution of singularities [69], using the De Concini–Procesi notion of wonderful models [23]. This argument allows one to simplify the formula for the multiplier ideals by eliminating a large number of redundant terms. My argument also remains in a finite-dimensional setting (unlike jet schemes, which are infinite-dimensional) and allows one to treat hyperplane arrangements with multiplicities.

Theorem 11 ([69]). *Let A be a hyperplane arrangement with nonnegative multiplicities in a vector space V and let $L(A)$ be the set of subspaces obtained as intersections of hyperplanes in A . For each $W \in L(A)$ let $r(W)$ be the codimension of W and let $s(W)$ be the sum of multiplicities of hyperplanes of A containing W . Let $\mathcal{G} \subset L(A)$ be any building set. Then the multiplier ideals of $I = I(A)$ are given by, for any $c \geq 0$,*

$$\mathcal{J}(I^c) = \bigcap_{W \in \mathcal{G}} I_W^{\lfloor c \cdot s(W) \rfloor - r(W) + 1},$$

where I_W is the ideal of W .

Mustaă's result is the case $\mathcal{G} = L(A) \setminus \{V\}$, but smaller building sets exist, and simplify the formula by reducing the number of ideals being intersected. For example, for the braid arrangement \mathcal{B}_n on \mathbb{C}^n , $|L(\mathcal{B}_n)|$ is the number of set partitions of $\{1, \dots, n\}$, which is super-exponential; while \mathcal{B}_n admits a building set (corresponding to so-called modular partitions) of size $2^n - n - 1$, i.e., exponential.

Future work. It remains unknown whether every member of a minimal building set is necessarily irredundant in the above formula, or whether a further reduction is possible.

2.2. Multiplier ideals of line arrangements. I computed the multiplier ideals of reduced unions of lines through the origin in \mathbb{C}^3 under certain hypotheses [76]. This is notable for being essentially the only computation of multiplier ideals carried out without special combinatorial or representation-theoretic structure. The computation was carried out using resolution of singularities. Thus, for example, for the ideal I_1 of three non-coplanar lines through the origin, $\mathcal{J}(I_1^c) = (1)$ for $0 \leq c < 3/2$, while for the ideal I_2 of three coplanar lines through the origin, $\mathcal{J}(I_2^c) = (1)$ for $0 \leq c < 5/3$. With my results, similar comparisons can be given in other examples: for instance, between 6 general lines through the origin and 6 lines through the origin lying on a quadratic cone.

Consideration of these line arrangements is motivated by the following example of Ein–Lazarsfeld–Smith [30]: If $Z \subset \mathbb{P}^2$ is a finite set of points, I is the homogeneous defining ideal of Z , $m > 0$ is a positive integer, and F is a homogeneous form vanishing to order at least

$2m$ at each point of Z , then $F \in I^m$. That is, the symbolic power $I^{(2m)}$ is contained in the ordinary power: $I^{(2m)} \subseteq I^m$. Despite the elementary nature of the statement, the only known proof is the one given by Ein–Lazarsfeld–Smith, using the asymptotic multiplier ideals of the line arrangement corresponding to the points Z . So computing the multiplier ideals of line arrangements is a natural first step toward a deeper understanding of such containments of symbolic powers. See for example [10, 9] where these are studied intensively and elementary proofs are given for the case that Z is a general set of points; these Ein–Lazarsfeld–Smith and Bocci–Harbourne papers sparked a great deal of (ongoing!) activity, see for example [39, 27, 28, 36, 8].

Future work. For arrangements of lines satisfying a certain hypothesis, a key part of the work in [76] is to show that certain exceptional divisors arising in the resolution of singularities are redundant. Compare [65, 78] where this phenomenon is studied for singularities of curves on a surface. I will study multiplier ideals of arbitrary line arrangements with an eye toward determining which exceptional divisors are redundant or irredundant.

2.3. Software for computing multiplier ideals. In theory it is algorithmic to compute multiplier ideals by computing a resolution of singularities of I followed by a sheaf pushforward. In practice it is more difficult, see [33].

Shibuta’s algorithm for computing Bernstein-Sato polynomials and multiplier ideals via Gröbner basis methods in Weyl algebras [62] (implemented by Shibuta in RISA/ASIR) was refined and implemented in the DMODULES library for MACAULAY2 by Berkesch and Leykin [4]. The DMODULES library can compute multiplier ideals and jumping numbers of arbitrary ideals, but due to the difficulty of the computations, can only handle modestly sized examples.

I have developed a new software package named MULTIPLIERIDEALS, see [71], that computes multiplier ideals of special ideals including monomial ideals, ideals of monomial curves, generic determinantal ideals, and hyperplane arrangements via combinatorial methods, using the NORMALIZ software and interface to MACAULAY2 by Bruns, et al [11, 12]. The combinatorial methods allow computations of somewhat larger examples than can be handled by general methods.

The MULTIPLIERIDEALS package also computes *jumping numbers*, those values of c at which $\mathcal{J}(I^c)$ changes, and the *log canonical threshold* $\text{lct}(I)$, the smallest positive jumping number.

The MULTIPLIERIDEALS package is available from my web site at <http://math.boisestate.edu/~zzeitler/math/MultiplierIdealsSoftware.php>. It has been submitted to the Journal of Software for Algebra and Geometry; upon acceptance it will be distributed as part of MACAULAY2.

Future work. I will continue to add to the software as more algorithms for multiplier ideals are developed. I will also work on applications of this software to statistical computations and other applications, as described in [80].

2.4. Monodromy of hyperplane arrangements. In joint work with Nero Budur and Mircea Mustața [15], we show that the Monodromy Conjecture holds for hyperplane arrangements and reduce a stronger version of the conjecture to a conjecture on Bernstein-Sato polynomials of hyperplane arrangements. We prove the latter conjecture for a class of hyperplane arrangements including generic arrangements and, using multiplier ideals, for

arrangements of “moderate type” (a certain monotonicity condition on multiplicities in the intersection lattice of the arrangement).

2.5. Asymptotic multiplier ideals of monomial ideals. The note [70], originally written as an appendix for lecture notes of Brian Harbourne [38], provides an exposition of asymptotic multiplier ideals and their application to the uniform bounds for symbolic powers developed by Ein–Lazarsfeld–Smith [30]. I show that a certain improvement of the Ein–Lazarsfeld–Smith bound found by Takagi–Yoshida [68] using characteristic p methods can also be obtained by the Ein–Lazarsfeld–Smith approach using asymptotic multiplier ideals:

Theorem 12 ([68]). *Let R be a regular local ring of equal characteristic 0, $I \subseteq R$ a reduced ideal, e be the greatest height of an associated prime of I , and ℓ an integer, $0 \leq \ell < \text{lct}(I^{(\bullet)})$ where $\text{lct}(I^{(\bullet)})$ is the log canonical threshold of the graded system of symbolic powers of I . Then $I^{(m)} \subseteq I^r$ whenever $m \geq er - \ell$. More generally, for any $k \geq 0$, $I^{(m)} \subseteq (I^{(k+1)})^r$ whenever $m \geq er + kr - \ell$.*

In addition I elaborate on an idea of Mustață [53] to compute asymptotic multiplier ideals of several families of graded sequences of monomial ideals, including in particular the sequence of symbolic powers of a reduced (squarefree) monomial ideal. These are among the very few nontrivial examples of asymptotic multiplier ideals of symbolic powers that have been computed to date. I also compute asymptotic multiplier ideals of graded systems of hyperplane arrangements.

3. EXPERIMENTS AT THE FRONTIERS OF REALITY IN SCHUBERT CALCULUS

I am a member of a team led by Frank Sottile to test a number of conjectures on reality in Schubert calculus by means of massive computational experiments.

Schubert calculus concerns the number of linear subspaces of \mathbb{C}^n having intersections of specified dimensions with some given flags (sequences of nested subspaces). For example, given 4 general planes in \mathbb{C}^4 , one finds that there are exactly 2 planes having at least 1-dimensional intersection with each — correspondingly, given 4 general lines in \mathbb{P}^3 , there are exactly 2 lines meeting each given line. When the given flags are *real*, the solution subspaces are real or conjugate pairs. “Reality in Schubert calculus” refers to the phenomenon that, for any given list of intersection conditions (called Schubert conditions), there exists some configuration of real flags for which the solution subspaces are all real. This was shown by Vakil [79] building on earlier work by Sottile [66]. Boris and Michael Shapiro conjectured, astonishingly, that an exceedingly simple recipe would produce flag configurations giving real solutions to all Schubert problems: namely, configurations of osculating flags to a fixed real moment curve. Interest in this conjecture, leading eventually to its proof by Mukhin–Tarasov–Varchenko [52], was spurred by massive computational experiments by Sottile and others, see for example [59].

I joined Sottile’s team which consisted of postdocs and graduate students working with Sottile. As a postdoc on the team, I mentored graduate students. We developed software to test variations of the Shapiro–Shapiro conjecture (Mukhin–Tarasov–Varchenko theorem), notably the *Secant Conjecture* which replaced osculating flags with secant flags spanned by points along the moment curve in disjoint intervals; the *Monotone Conjecture* which generalized the Shapiro–Shapiro conjecture to the setting of flag manifolds instead of Grassmannians; and the *Monotone Secant Conjecture*. These all remain open. In addition to providing overwhelming evidence for all three conjectures, our experiments explored new

territory and uncovered new phenomena by including computations of secant flags along non-disjoint intervals, and non-monotone configurations.

The results of our experiments are reported in [34, 40]. The running of the experiments is described in [41]. All the data generated by the experiments is publicly accessible through the project web site, <http://www.math.tamu.edu/~secant/>.

The computations were carried out primarily using SINGULAR and MAPLE, as well as MACAULAY2, CoCoA, and SAGE. We used PERL and batch scheduling software to automate the computation of billions of examples. Results were stored in a database using MYSQL and displayed on dynamically created web pages using PHP. I was involved in the development of every part of the codebase.

4. ARRANGEMENT, COMBINATORIAL, AND DETERMINANTAL PROBLEMS

My study of multiplier ideals of line arrangements led me to look into arrangements of points, hyperplanes, and more generally arrangements of linear subspaces. As points in the plane are defined by the maximal minors of a Hilbert–Burch matrix, I also became interested in certain questions about determinantal ideals. My activities in these areas include the following.

4.1. Hilbert functions of fat point schemes in the plane. This is joint work with Susan Cooper and Brian Harbourne [22]. Let A be a fat point scheme in the plane, $A = m_1P_1 + \cdots + m_nP_n$, and suppose we are given certain subsets $S_1, \dots, S_k \subseteq \{P_1, \dots, P_n\}$ such that for each i , there exists a line L_i containing the points in S_i and no other points of A ; but we do not assume knowledge of the lines L_i , the positions of the points along each L_i , actual coordinates of the P_j , etc. We also do not assume that the given list of S_i is complete, i.e., there may be additional subsets of collinear points, but we are limited to just the given S_i .

From this limited information we describe a simple recursive reduction procedure as follows: at each step, choose a subset S_i , record the total of multiplicities of points in S_i , then decrease each of those multiplicities by 1, discarding any point whose multiplicity reaches zero. The output is a vector of nonnegative integers, the totals of multiplicities recorded at each step. This is called a *reduction vector*.

Also for any vector d of nonnegative integers we define functions $f_d, F_d : \mathbb{Z} \rightarrow \mathbb{Z}$, and show:

Theorem 13 ([22]). *If d is a reduction vector of A then the Hilbert function h_A satisfies $f_d \leq h_A \leq F_d$. Furthermore, if d contains no zero entries and is non-increasing, then $f_d = F_d$, unless d contains a subsequence of the form (\dots, a, a, a, \dots) or $(\dots, a, a, a + 1, a + 2, \dots, a + k - 1, a + k, a + k, \dots)$, that is, three equal entries, or consecutive entries with the first and last repeated.*

Conversely if $f_d = F_d$ then d satisfies the condition, i.e., avoids the indicated subsequences. Both f_d and F_d are defined recursively but are also given explicit combinatorial descriptions. Interestingly, “greedy” choices (i.e., choosing at each step a subset S_i to have the largest possible sum of multiplicities) do not necessarily result in the best possible bounds, as we show by example.

In the case d avoids the indicated subsequences, so that $f_d = F_d$ and h_A is uniquely determined, we also give upper and lower bounds for the graded Betti numbers of the ideal I_A , and we show that those bounds coincide precisely when the reduction vector d is strictly decreasing.

We compute the Hilbert functions and graded Betti numbers for a number of interesting examples, such as star configurations with multiplicities.

4.2. Complete bipartite subspace arrangements. This is joint work with Douglas A. Torrance [74]. We consider arrangements of codimension 2 subspaces about which one has only the following partial information: the simple graph with one vertex for each subspace and links whenever the subspaces have codimension 3 intersection. This graph does not typically determine the Castelnuovo–Mumford regularity of the (ideal of the) arrangement, or whether the arrangement is arithmetically Cohen–Macaulay. However we show that in case the graph is a complete bipartite graph of type (a, b) with $a \leq b \leq 2$ or $2 \leq a \leq b \leq 3$ or $3 \leq a \leq b$, the regularity of the ideal is uniquely determined, in fact equal to $\max(a + 1, b)$; and the arrangement A is arithmetically Cohen–Macaulay if and only if $b = a$ or $b = a + 1$.

4.3. Decompositions of ideals of minors meeting a submatrix. This is joint work with Kent M. Neuberger [56]. We give primary decompositions of certain ideals generated by subsets of minors or Pfaffians of a generic matrix. First, let $X = (x_{i,j})$ be a generic matrix. It is well known that the ideal $I_t(X)$ of t -minors of X is prime. Now consider the ideal generated by those t -minors that involve at least t_i columns in the first c_i columns of X , for each i , for some given values t_i, c_i ; and one can similarly impose row conditions. These ideals arise when, for example, one considers replacement of the $x_{i,j}$ by homogeneous forms of degree $b_j - a_i$; then the low-degree part of the resulting determinantal ideal is described by these sorts of row and column conditions. We give an explicit primary decomposition of these determinantal ideals, using the theory of algebras with straightening law. This generalizes one of the results of [1] concerning the ideal generated by minors that contain a submatrix, i.e., involve all c_i of the first c_i columns of X (and all the rows).

We find similar results for minors in a generic symmetric matrix and Pfaffians in a generic skew-symmetric matrix.

5. OTHER

Not discussed here are my papers on the following:

- on nef cone volumes of generalized Del Pezzo surfaces (joint work with Ulrich Derenthal and Michael Joyce) [24];
- a report on recent developments and open problems in linear series [3] (I contributed a section on bounds for symbolic powers);
- on arithmetic forms of toric varieties, i.e., with nonstandard structures over nonclosed fields (joint work with Javier Elizondo, Paulo Lima-Filho, and Frank Sottile) [32];
- generalizing Chris Hammond’s topological criterion for schlichtness of the domain of holomorphy of a function [72];
- on intersections of curves through a set of points in the plane [77].

Research Impact: Two problems or conjectures that I suggested have led to papers by other researchers.

- A problem which I suggested to Dan Erman was used as the basis for a successful research program by a group of graduate students at the MSRI Summer Graduate Workshop in Commutative Algebra in summer 2011. The problem was to find the Boij–Söderberg decomposition for the Betti table of a complete intersection. The group of graduate students under Dan’s direction found partial results toward this

problem, leading to the recent paper *Non-simplicial decompositions of Betti diagrams of complete intersections* by Courtney Gibbons, Jack Jeffries, Sarah Mayes, Claudiu Raicu, Branden Stone, and Bryan White (arXiv:1301.3441 [math.AC]) (to appear in Journal of Commutative Algebra). Understanding this (very) special case is a first step toward understanding more interesting cases, and suggests more questions for future investigation.

- A conjecture which I shared with A. Iarrobino was solved by his recent graduate student Masoumeh (Sepideh) Shafiei, who determined the apolar annihilating ideals of the generic determinant, permanent, Pfaffian, Hafnian, symmetric determinant, and more. These results were the basis of her Ph.D. dissertation and led to her recent papers *Apolarity for determinants and permanents of generic matrices* (arXiv:1212.0515) (to appear in Journal of Commutative Algebra) and *Apolarity for determinants and permanents of generic symmetric matrices* (arXiv:1303.1860). They have already found application in my ongoing research.

Other problems. I am interested in studying the sequences of regularities that occur for exact sequences of graded modules, and the “Betti tensor” obtained by stacking the Betti tables of graded modules in an exact sequence, generalizing Boij–Söderberg theory.

I will study jet schemes of monomial ideals and subspace arrangements, building on work of Mustața, Goward–Smith, and C. Yuen.

I would like to extend Jeremy Marin’s slope varieties and picture varieties to higher dimension and codimension and other settings such as tropical geometry.

REFERENCES

1. J. F. Andrade and A. Simis, *On ideals of minors fixing a submatrix*, J. Algebra **102** (1986), no. 1, 246–259. MR 853243 (87j:13028)
2. Edoardo Ballico and Alessandro De Paris, *Generic power sum decompositions and bounds for the Waring rank*, arXiv:1312.3494 [math.AG], Dec 2013.
3. Thomas Bauer, Cristiano Bocci, Susan Cooper, Sandra Di Rocco, Marcin Dumnicki, Brian Harbourne, Kelly Jabbusch, Andreas Leopold Knutsen, Alex Küronya, Rick Miranda, Joaquim Roé, Hal Schenck, Tomasz Szemberg, and Zach Teitler, *Recent developments and open problems in linear series*, Contributions to algebraic geometry, EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2012, pp. 93–140. MR 2976940
4. Christine Berkesch and Anton Leykin, *Algorithms for Bernstein–Sato polynomials and multiplier ideals*, ISSAC (Wolfram Koepf, ed.), ACM, 2010, pp. 99–106.
5. A. Białynicki-Birula and A. Schinzel, *Representations of multivariate polynomials by sums of univariate polynomials in linear forms*, Colloq. Math. **112** (2008), no. 2, 201–233. MR 2383331 (2009b:12006)
6. Caucher Birkar, Paolo Cascini, Christopher D. Hacon, and James M^cKernan, *Existence of minimal models for varieties of log general type*, J. Amer. Math. Soc. **23** (2010), no. 2, 405–468. MR 2601039
7. Grigoriy Blekherman and Zach Teitler, *On Maximum, Typical, and Generic Ranks*, arXiv:1402.2371 [math.AG], Feb 2014.
8. Cristiano Bocci, Susan M. Cooper, and Brian Harbourne, *Containment results for ideals of various configurations of points in \mathbf{P}^N* , J. Pure Appl. Algebra **218** (2014), no. 1, 65–75. MR 3120609
9. Cristiano Bocci and Brian Harbourne, *Comparing powers and symbolic powers of ideals*, J. Algebraic Geom. **19** (2010), no. 3, 399–417. MR 2629595 (2011d:13021)
10. ———, *The resurgence of ideals of points and the containment problem*, Proc. Amer. Math. Soc. **138** (2010), no. 4, 1175–1190. MR 2578512 (2011f:14010)
11. Winfried Bruns and Bogdan Ichim, *Normaliz: algorithms for affine monoids and rational cones*, J. Algebra **324** (2010), no. 5, 1098–1113. MR 2659215

12. Winfried Bruns and Gesa Kämpf, *A Macaulay2 interface for Normaliz*, J. Softw. Algebra Geom. **2** (2010), 15–19. MR 2881130
13. Weronika Buczyńska, Jarosław Buczyński, Johannes Kleppe, and Zach Teitler, *Apolarity and direct sum decomposability of polynomials*, arXiv:1307.3314 [math.AG], Jul 2013.
14. Weronika Buczyńska, Jarosław Buczyński, and Zach Teitler, *Waring decompositions of monomials*, J. Algebra **378** (2013), 45–57. MR 3017012
15. Nero Budur, Mircea Mustața, and Zach Teitler, *The monodromy conjecture for hyperplane arrangements*, Geom. Dedicata **153** (2011), 131–137. MR 2819667 (2012i:32035)
16. Nero Budur and Morihiko Saito, *Multiplier ideals, V -filtration, and spectrum*, J. Algebraic Geom. **14** (2005), no. 2, 269–282. MR 2123230 (2006g:14012)
17. Enrico Carlini, *Codimension one decompositions and Chow varieties*, Projective varieties with unexpected properties, Walter de Gruyter GmbH & Co. KG, Berlin, 2005, pp. 67–79. MR 2202247 (2007f:14056)
18. ———, *Binary decompositions and varieties of sums of binaries*, J. Pure Appl. Algebra **204** (2006), no. 2, 380–388. MR 2184818 (2006j:14070)
19. Enrico Carlini, Maria Virginia Catalisano, and Luca Chiantini, *Progress on the symmetric Strassen conjecture*, arXiv:1405.3721 [math.AG], May 2014.
20. Enrico Carlini, Maria Virginia Catalisano, and Anthony V. Geramita, *The solution to the Waring problem for monomials and the sum of coprime monomials*, J. Algebra **370** (2012), 5–14.
21. Pierre Comon, Gene Golub, Lek-Heng Lim, and Bernard Mourrain, *Symmetric tensors and symmetric tensor rank*, SIAM J. Matrix Anal. Appl. **30** (2008), no. 3, 1254–1279. MR 2447451 (2009i:15039)
22. Susan Cooper, Brian Harbourne, and Zach Teitler, *Combinatorial bounds on hilbert functions of fat points in projective space*, J. Pure Appl. Algebra **215** (2011), no. 9, 2165–2179.
23. C. De Concini and C. Procesi, *Wonderful models of subspace arrangements*, Selecta Math. (N.S.) **1** (1995), no. 3, 459–494. MR MR1366622 (97k:14013)
24. Ulrich Derenthal, Michael Joyce, and Zachariah Teitler, *The nef cone volume of generalized Del Pezzo surfaces*, Algebra Number Theory **2** (2008), no. 2, 157–182. MR MR2377367 (2009b:14069)
25. Harm Derksen and Zach Teitler, *Lower bound for ranks of invariant forms*, arXiv:1409.0061 [math.AG], Aug 2014.
26. Mathias Drton, Bernd Sturmfels, and Seth Sullivant, *Lectures on algebraic statistics*, Oberwolfach Seminars, vol. 39, Birkhäuser Verlag, Basel, 2009. MR 2723140 (2012d:62004)
27. Marcin Dumnicki, Brian Harbourne, Tomasz Szemberg, and Halszka Tutaj-Gasińska, *Linear subspaces, symbolic powers and Nagata type conjectures*, Adv. Math. **252** (2014), 471–491. MR 3144238
28. Marcin Dumnicki, Tomasz Szemberg, and Halszka Tutaj-Gasińska, *Counterexamples to the $I^{(3)} \subset I^2$ containment*, J. Algebra **393** (2013), 24–29. MR 3090054
29. Lawrence Ein, Robert Lazarsfeld, and Mircea Mustața, *Contact loci in arc spaces*, Compos. Math. **140** (2004), no. 5, 1229–1244. MR 2081163 (2005f:14006)
30. Lawrence Ein, Robert Lazarsfeld, and Karen E. Smith, *Uniform bounds and symbolic powers on smooth varieties*, Invent. Math. **144** (2001), no. 2, 241–252. MR MR1826369 (2002b:13001)
31. Lawrence Ein and Mircea Mustața, *Invariants of singularities of pairs*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, 2006, pp. 583–602. MR MR2275611 (2007m:14050)
32. E. Javier Elizondo, Paulo Lima-Filho, Frank Sottile, and Zach Teitler, *Arithmetic toric varieties*, Math. Nachr. **287** (2014), no. 2-3, 216–241. MR 3163576
33. Anne Frühbis-Krüger, *Desingularization in computational applications and experiments*, arXiv:1301.3709 [math.AG], Jan 2013.
34. Luis D. García-Puente, Nickolas Hein, Christopher Hillar, Abraham Martín del Campo, James Ruffo, Frank Sottile, and Zach Teitler, *The Secant Conjecture in the Real Schubert Calculus*, Experimental Mathematics **21** (2012), no. 3, 252–265.
35. Anthony V. Geramita, *Inverse systems of fat points: Waring’s problem, secant varieties of Veronese varieties and parameter spaces for Gorenstein ideals*, The Curves Seminar at Queen’s, Vol. X (Kingston, ON, 1995), Queen’s Papers in Pure and Appl. Math., vol. 102, Queen’s Univ., Kingston, ON, 1996, pp. 2–114.
36. Elena Guardo, Brian Harbourne, and Adam Van Tuyl, *Fat lines in \mathbb{P}^3 : powers versus symbolic powers*, J. Algebra **390** (2013), 221–230. MR 3072120

37. Christopher D. Hacon and James M^cKernan, *Extension theorems and the existence of flips*, Flips for 3-folds and 4-folds, Oxford Lecture Ser. Math. Appl., vol. 35, Oxford Univ. Press, Oxford, 2007, pp. 76–110. MR MR2359343
38. Brian Harbourne, *Global aspects of the geometry of surfaces*, Ann. Univ. Paedagog. Crac. Stud. Math. **9** (2010), 5–41. MR 2608654 (2011g:14012)
39. Brian Harbourne and Craig Huneke, *Are symbolic powers highly evolved?*, J. Ramanujan Math. Soc. **28A** (2013), 247–266. MR 3115195
40. Nickolas Hein, Christopher J. Hillar, Abraham Martín del Campo, Frank Sottile, and Zach Teitler, *The monotone secant conjecture in the real Schubert calculus*, arXiv:1109.3436 [math.AG], Jun 2014.
41. Christopher Hillar, Luís García-Puente, Abraham Martín del Campo, James Ruffo, Zach Teitler, Stephen L. Johnson, and Frank Sottile, *Experimentation at the Frontiers of Reality in Schubert Calculus*, Gems in Experimental Mathematics (Tewodros Amdeberhan, Luis A. Medina, and Victor H. Moll, eds.), Contemp. Math., vol. 517, AMS, 2010, pp. 365–380.
42. Erik Holmes, Paul Plummer, Jeremy Siegert, and Zach Teitler, *Maximum Waring ranks of monomials*, arXiv:1309.7834 [math.AG], Apr 2014.
43. Anthony Iarrobino and Vassil Kanev, *Power sums, Gorenstein algebras, and determinantal loci*, Lecture Notes in Mathematics, vol. 1721, Springer-Verlag, Berlin, 1999, Appendix C by Iarrobino and Steven L. Kleiman. MR 1735271 (2001d:14056)
44. Joachim Jelisiejew, *An upper bound for the Waring rank of a form*, Arch. Math. (Basel) **102** (2014), no. 4, 329–336. MR 3196960
45. J. J. Kohn, *Sufficient conditions for subellipticity on weakly pseudo-convex domains*, Proc. Nat. Acad. Sci. U.S.A. **74** (1977), no. 6, 2214–2216. MR 0466635 (57 #6512)
46. ———, *Subellipticity of the $\bar{\partial}$ -Neumann problem on pseudo-convex domains: sufficient conditions*, Acta Math. **142** (1979), no. 1-2, 79–122. MR 512213 (80d:32020)
47. Tamara G. Kolda and Brett W. Bader, *Tensor decompositions and applications*, SIAM Rev. **51** (2009), no. 3, 455–500. MR 2535056 (2010j:15027)
48. J. M. Landsberg, *Tensors: geometry and applications*, Graduate Studies in Mathematics, vol. 128, American Mathematical Society, Providence, RI, 2012. MR 2865915
49. J.M. Landsberg and Zach Teitler, *On the ranks and border ranks of symmetric tensors*, Found. Comp. Math. **10** (2010), no. 3, 339–366.
50. Robert Lazarsfeld, *Positivity in algebraic geometry. II*, Ergebnisse der Mathematik., vol. 49, Springer-Verlag, Berlin, 2004, Positivity for vector bundles, and multiplier ideals. MR MR2095472
51. Yongnam Lee, *Chow stability criterion in terms of log canonical threshold*, J. Korean Math. Soc. **45** (2008), no. 2, 467–477. MR 2389549 (2009b:14091)
52. Evgeny Mukhin, Vitaly Tarasov, and Alexander Varchenko, *The B. and M. Shapiro conjecture in real algebraic geometry and the Bethe ansatz*, Ann. of Math. (2) **170** (2009), no. 2, 863–881. MR 2552110 (2011b:17065)
53. Mircea Mustața, *On multiplicities of graded sequences of ideals*, J. Algebra **256** (2002), no. 1, 229–249. MR MR1936888 (2003k:13030)
54. ———, *Multiplier ideals of hyperplane arrangements*, Trans. Amer. Math. Soc. **358** (2006), 5015–5023.
55. Alan Michael Nadel, *Multiplier ideal sheaves and Kähler-Einstein metrics of positive scalar curvature*, Ann. of Math. (2) **132** (1990), no. 3, 549–596. MR 1078269 (92d:32038)
56. Kent M. Neuerburg and Zach Teitler, *Decompositions of ideals of minors meeting a submatrix*, arXiv:1406.6426 [math.AC], Jun 2014.
57. Wenbo Niu, *Singularities of generic linkage of algebraic varieties*, arXiv:1207.1082 [math.AG], Jul 2012.
58. Kristian Ranestad and Frank-Olaf Schreyer, *On the rank of a symmetric form*, J. Algebra **346** (2011), 340–342. MR 2842085
59. Jim Ruffo, Yuval Sivan, Evgenia Soprunova, and Frank Sottile, *Experimentation and conjectures in the real Schubert calculus for flag manifolds*, Experiment. Math. **15** (2006), no. 2, 199–221. MR MR2253007 (2007g:14066)
60. Masoumeh Sepideh Shafiei, *Apolarity for determinants and permanents of generic symmetric matrices*, arXiv:1303.1860 [math.AC], Mar 2013.

61. ———, *Apolarity for determinants and permanents of generic matrices*, To appear in Journal of Commutative Algebra, 2014.
62. Takafumi Shibuta, *Algorithms for computing multiplier ideals*, Journal of Pure and Applied Algebra **215** (2011), no. 12, 2829–2842.
63. Yum-Tong Siu, *Invariance of plurigenera*, Invent. Math. **134** (1998), no. 3, 661–673. MR MR1660941 (99i:32035)
64. Karen E. Smith, *The multiplier ideal is a universal test ideal*, Comm. Algebra **28** (2000), no. 12, 5915–5929, Special issue in honor of Robin Hartshorne. MR 1808611 (2002d:13008)
65. Karen E. Smith and Howard M. Thompson, *Irrelevant exceptional divisors for curves on a smooth surface*, Algebra, geometry and their interactions, Contemp. Math., vol. 448, Amer. Math. Soc., Providence, RI, 2007, pp. 245–254. MR MR2389246
66. Frank Sottile, *Some real and unreal enumerative geometry for flag manifolds*, Michigan Math. J. **48** (2000), 573–592, Dedicated to William Fulton on the occasion of his 60th birthday. MR MR1786506 (2002d:14085)
67. J.J. Sylvester, *An essay on canonical forms, supplement to a sketch of a memoir on elimination, transformation and canonical forms*, originally published by George Bell, Fleet Street, London, 1851. Paper 34 in *Mathematical Papers*, Vol. 1, Chelsea, New York, 1973, originally published by Cambridge University Press in 1904., 1851.
68. Shunsuke Takagi and Ken-ichi Yoshida, *Generalized test ideals and symbolic powers*, Michigan Math. J. **57** (2008), 711–724.
69. Zach Teitler, *A note on Mustață’s computation of multiplier ideals of hyperplane arrangements*, Proc. Amer. Math. Soc. **136** (2008), no. 5, 1575–1579. MR 2373586 (2008k:14005)
70. ———, *Bounding symbolic powers via asymptotic multiplier ideals*, Ann. Univ. Paedagog. Crac. Stud. Math. **8** (2009), 67–77. MR 2591736
71. ———, *Software for multiplier ideals*, arXiv:1305.4435 [math.AG], May 2013.
72. ———, *Topological criteria for schlichtness*, Proc. Edinb. Math. Soc. (2) **56** (2013), no. 2, 637–640. MR 3056664
73. ———, *Geometric lower bounds for generalized ranks*, arXiv:1406.5145 [math.AG], Jun 2014.
74. Zach Teitler and Douglas A. Torrance, *Castelnuovo-Mumford regularity and arithmetic Cohen-Macaulayness of complete bipartite subspace arrangements*, JPAA (2014), DOI: 10.1016/j.jpaa.2014.07.027.
75. Zach Teitler and Alex Woo, *Apolarity and reflection groups*, J. Alg. Comb. (2014), DOI: 10.1007/s10801-014-0539-0.
76. Zachariah C. Teitler, *Multiplier ideals of general line arrangements in \mathbb{C}^3* , Comm. Algebra **35** (2007), no. 6, 1902–1913.
77. ———, *On the intersection of the curves through a set of points in \mathbb{P}^2* , JPAA **209** (2007), no. 2, 571–581.
78. Kevin Tucker, *Jumping numbers on algebraic surfaces with rational singularities*, Trans. Amer. Math. Soc. **362** (2010), no. 6, 3223–3241. MR 2592954 (2011c:14106)
79. Ravi Vakil, *Schubert induction*, Ann. of Math. (2) **164** (2006), no. 2, 489–512. MR MR2247966 (2007j:14082)
80. Sumio Watanabe, *Algebraic geometry and statistical learning theory*, Cambridge Monographs on Applied and Computational Mathematics, vol. 25, Cambridge University Press, Cambridge, 2009. MR 2554932 (2011g:62185)
81. Youngho Woo, *Some cases on Strassen additive conjecture*, arXiv:1406.2213 [math.AG], Jun 2014.
82. Piotr Zwiernik, *An asymptotic behaviour of the marginal likelihood for general Markov models*, J. Mach. Learn. Res. **12** (2011), 3283–3310. MR 2877601 (2012m:62248)